

QUATERNIONIC QUASIDETERMINANTS AND DETERMINANTS

ISRAEL GELFAND, VLADIMIR RETAKH, ROBERT LEE WILSON

INTRODUCTION

Quasideterminants of noncommutative matrices introduced in [GR, GR1] have proved to be a powerful tool in basic problems of noncommutative algebra and geometry (see [GR, GR1-GR4, GKLLRT, GV, EGR, EGR1, ER, KL, KLT, LST, Mo, Mo1, P, RS, RRV, Rsh, Sch]). In general, the quasideterminants of matrix $A = (a_{ij})$ are rational functions in (a_{ij}) 's. The minimal number of successive inversions required to express an rational function is called the height of this function. The "inversion height" is an important invariant showing a degree of "noncommutativity".

In general, the height of the quasideterminants of matrices of order n equals $n - 1$ (see [Re]). Quasideterminants are most useful when, as in commutative case, their height is less or equal to 1. Such examples include quantum matrices and their generalizations, matrices of differential operators, etc. (see [GR, GR1-GR4, ER]).

Quasideterminants of quaternionic matrices $A = (a_{ij})$ provide a closely related example. They can be written as a ratio of polynomials in a_{ij} 's and their conjugates. In fact, our Theorem 3.3 shows that any quasideterminant of A is a sum of monomials in the a_{ij} 's and the $\bar{a}_{ij}$'s with real coefficients.

The theory of quasideterminants leads to a natural definition of determinants of square matrices over noncommutative rings.

There is a long history of attempts to develop such a theory. These works have resulted in a number of useful generalizations of determinants to special classes of noncommutative rings, e.g. [B, C, Ca, Di, Dy1, F, H, KS, O, R, SK, W] a.o. to noncommutative determinants. In all these cases noncommutative determinants can be expressed as products of quasideterminants. It shows that quasideterminants tend to be a more fundamental notion in noncommutative algebra.

In particular, studies of quaternionic linear algebra initiated by Hamilton has resulted in notions of determinants due to Study and Moore and their applications [Ad, Al, As, Dy, Dy1]. We show here that these definitions may be obtained from a general procedure as products of quasideterminants.

A good exposition of quaternionic determinants can be found in [As].

The results of this paper can be easily generalized to a large variety of algebras with anti-involutions.

We would like to thank S. Alesker, A. Barvinok, and I. Frenkel. Their questions about relations between quasideterminants and determinants were very stimulating.

The second author was partially supported by the National Science Foundation.

I. PRELIMINARY

1.1. Quaternions.

The algebra of quaternions $\mathbb{H}$ is a unital $\mathbb{R}$ -algebra with generators $\mathbf{i}, \mathbf{j}, \mathbf{k}$ such that

$$\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = -1, \quad \mathbf{ij} = \mathbf{k} = -\mathbf{ji}, \quad \mathbf{jk} = \mathbf{i} = -\mathbf{kj}, \quad \mathbf{ki} = \mathbf{j} = -\mathbf{ik}.$$

Elements $h \in \mathbb{H}$ can be written as $h = a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}$ where $a, b, c, d \in \mathbb{R}$. Conjugation $h \mapsto \bar{h}$ in $\mathbb{H}$ is defined by $\bar{h} = a - b\mathbf{i} - c\mathbf{j} - d\mathbf{k}$ and satisfies $\overline{xy} = \bar{y}\bar{x}$. It follows that $h\bar{h} = \bar{h}h$.

Define the norm $\nu(h)$ of a quaternion h by setting $\nu(h) = h\bar{h}$. The norm $\nu(h)$ is a non-negative real number and $\nu(h) = 0$ if and only if $h = 0$. By definition, $|a| = \sqrt{\nu(a)}$. It follows that h^{-1} is equal to $\bar{h}$ up to a central coefficient:

$$h^{-1} = \frac{1}{\nu(a)} \bar{h}.$$

The field of real numbers $\mathbb{R}$ is canonically identified with a subfield in $\mathbb{H}$ via embedding $a \mapsto a + 0\mathbf{i} + 0\mathbf{j} + 0\mathbf{k}$. Then $h = \bar{h}$ if and only if $h \in \mathbb{R}$. The center of $\mathbb{H}$ coincides with $\mathbb{R}$.

An embedding of the field of complex numbers $\mathbb{C}$ into $\mathbb{H}$ is defined by an image of $\mathbf{i} \in \mathbb{C}$. We chose the embedding given by $a + b\mathbf{i} \mapsto a + b\mathbf{i} + 0\mathbf{j} + 0\mathbf{k}$, where $a, b \in \mathbb{R}$. The embedding $\mathbb{C}$ into $\mathbb{H}$ agrees with the conjugation.

Under these identifications every quaternionic number h can be written as $h = \alpha + \mathbf{j}\beta$ where $\alpha, \beta \in \mathbb{C}$.

Let $M(n, F)$ be the algebra of $n \times n$ -matrices over a field F . Define a map $\theta : \mathbb{H} \rightarrow M(2, \mathbb{C})$ by setting

$$\theta(h) = \begin{pmatrix} \alpha & -\bar{\beta} \\ \beta & \bar{\alpha} \end{pmatrix}.$$

It is well-known that θ is a homomorphism of $\mathbb{R}$ -algebras. Also, $\det \theta(h) = \nu(h)$.

If matrix $A = (a_{ij})$, $i, j = 1, \dots, n$ is a quaternionic matrix, define its Hermitian dual matrix to be $A^* = (b_{ij})$ where $b_{ij} = \bar{a}_{j,i}$ for all i, j . A is a Hermitian matrix if $A = A^*$.

1.2. Quasideterminants.

1.1.2. Definition of quasideterminants.

The definition of quasideterminants and their main properties were given in [GR. GR1-GR4].

Let $A = (a_{ij}), i \in I, j \in J$ be a matrix over a ring R . Construct the following submatrices of A : submatrix $A^{ij}, i \in I, j \in J$ obtained from A by deleting its i -th row and j -th column; row submatrix r_i^j obtained from i -th row of A by deleting the element a_{ij} ; column submatrix c_j^i obtained from j -th column of A by deleting the element a_{ij} .

Definition 1.1. *Let A be a square matrix. We say that the quasideterminant $|A|_{ij}$ is defined if the submatrix A^{ij} is invertible over the ring R . In this case*

$$(1.1) \quad |A|_{ij} = a_{ij} - r_i^j (A^{ij})^{-1} c_j^i.$$

We set $|A|_{ij} = a_{ij}$ if $I = \{i\}, J = \{j\}$.

If inverses to the quasideterminants $|A^{ij}|_{pq}, p \neq i, q \neq j$, are defined then

$$(1.2) \quad |A|_{ij} = a_{ij} - \sum a_{iq} |A^{ij}|_{pq}^{-1} a_{pj}.$$

Here the sum is taken over all $p \in I \setminus \{i\}, q \in J \setminus \{j\}$.

If A is an $n \times n$ -matrix there exist up to n^2 quasideterminants of A .

If A is a square matrix we call a quasideterminant of any $k \times k$ -submatrix a k -quasiminor of A .

Examples 1.2.

1) For a matrix $A = (a_{ij}), i, j = 1, 2$ there exist four quasideterminants if the corresponding entries are invertible

$$\begin{aligned} |A|_{11} &= a_{11} - a_{12} \cdot a_{22}^{-1} \cdot a_{21}, & |A|_{12} &= a_{12} - a_{11} \cdot a_{21}^{-1} \cdot a_{22}, \\ |A|_{21} &= a_{21} - a_{22} \cdot a_{12}^{-1} \cdot a_{11}, & |A|_{22} &= a_{22} - a_{21} \cdot a_{11}^{-1} \cdot a_{12}. \end{aligned}$$

If an entry, say a_{22} is not invertible then the quasideterminant $|A|_{11}$ does not exist.

2) For a matrix $A = (a_{ij}), i, j = 1, 2, 3$ there exist up to 9 quasideterminants but we will write here only

$$\begin{aligned} |A|_{11} &= a_{11} - a_{12}(a_{22} - a_{23}a_{33}^{-1}a_{32})^{-1}a_{21} & -a_{12}(a_{32} - a_{33} \cdot a_{23}^{-1}a_{22})^{-1}a_{31} \\ & -a_{13}(a_{23} - a_{22}a_{32}^{-1}a_{33})^{-1}a_{21} & -a_{13}(a_{33} - a_{32} \cdot a_{22}^{-1}a_{23})^{-1}a_{31} \end{aligned}$$

provided all inverses are defined.

Remark. If each a_{ij} is an invertible morphism $V_j \rightarrow V_i$ in an additive category then the quasideterminant $|A|_{pq}$ is also a morphism from the object V_q to the object V_p .

Example 1.3.

Quasideterminants are not generalizations of determinants over a commutative ring but generalizations of the ratio of two determinants. If in formulas (1.1), or (1.2) the variables a_{ij} commute then

$$|A|_{pq} = (-1)^{p+q} \frac{\det A}{\det A^{pq}}.$$

As a general rule, quasideterminants are rational functions in their entries but not polynomials.

The following theorem was conjectured by I. Gelfand and V. Retakh, and proved by Reutenauer [Re].

Theorem 1.4. *Quasideterminants of the $n \times n$ -matrix with formal entries have the inversion height $n - 1$ over the free skew-field generated by its entries.*

1.3. General properties of quasideterminants**Elementary properties of quasideterminants.**

- i) The quasideterminant $|A|_{pq}$ does not depend on the permutation of rows and columns in the matrix A if the p -th row and the q -th column are not changed;
- ii) *The multiplication of rows and columns.* Let the matrix B be constructed from the matrix A by multiplication of its i -th row by a scalar λ from the left. Then

$$|B|_{kj} = \begin{cases} \lambda |A|_{ij} & \text{if } k = i \\ |A|_{kj} & \text{if } k \neq i \text{ and } \lambda \text{ is invertible.} \end{cases}$$

Let the matrix C be constructed from the matrix A by multiplication of its j -th column by a scalar μ from the right. Then

$$|C|_{i\ell} = \begin{cases} |A|_{ij}\mu & \text{if } \ell = j \\ |A|_{i\ell} & \text{if } \ell \neq j \text{ and } \mu \text{ is invertible.} \end{cases}$$

- iii) *The addition of rows and columns.* Let the matrix B be constructed by adding to the p -th row of the matrix A its k -th row multiplied by a scalar λ from the left, where $k \neq p$. Then

$$|A|_{ij} = |B|_{ij}, \quad i = 1, \dots, k-1, k+1, \dots, n, j = 1, \dots, n.$$

Let the matrix C be constructed by addition to the q -th column of the matrix A its ℓ -th column multiplied by a scalar λ from the right, where $\ell \neq q$. Then

$$|A|_{ij} = |C|_{ij}, \quad i = 1, \dots, n, j = 1, \dots, \ell-1, \ell+1, \dots, n.$$

Homological relations. The ratio of two quasideterminants of the same square matrix is the ratio of two quasiminors. This is a consequence of the following *homological* relations [GR, GR1]:

Theorem 1.5.

a) *Row homological relations:*

$$-|A|_{ij} \cdot |A^{i\ell}|_{sj}^{-1} = |A|_{i\ell} \cdot |A^{ij}|_{s\ell}^{-1} \quad \forall s \neq i$$

b) *Column homological relations:*

$$-|A^{kj}|_{it}^{-1} \cdot |A|_{ij} = |A^{ij}|_{kt}^{-1} \cdot |A|_{kj} \quad \forall r \neq j$$

Heredity. Let $A = (a_{ij}), i, j = 1, \dots, n$, be a matrix and

$$A = \begin{pmatrix} A_{11} & \dots & A_{1s} \\ A_{s1} & \dots & A_{ss} \end{pmatrix}$$

be a block decomposition of A . Denote by $\tilde{A} = (A_{ij}), i, j = 1, \dots, s$, the matrix with A_{ij} 's as entries. Suppose that

$$A_{pq} = \begin{pmatrix} a_{k\ell} & \dots & a_{k,\ell+m} \\ & \dots & \\ a_{k+m,\ell} & \dots & a_{k+m,\ell+m} \end{pmatrix}.$$

is a square matrix and that $|\tilde{A}|_{pq}$ is defined.

Theorem 1.6.

$$|A|_{k'\ell'} = |\tilde{A}|_{pq}|_{k'\ell'} \text{ for } k' = k, \dots, k+m, \ell' = \ell, \dots, \ell+m.$$

In particular, let $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}$ be a block decomposition of $A = (a_{ij}), i, j = 1, \dots, n$. Let A_{11} be a $k \times k$ -matrix and let the matrix A_{22} be invertible.

Then $|A|_{ij} = |A_{11} - A_{12}A_{22}^{-1} \cdot A_{21}|_{ij}$ for $i, j = 1, \dots, k$.

In other words, the quasideterminant $|A|_{ij}$ of an $n \times n$ matrix can be computed in two steps: first, consider the quasideterminant $|\tilde{A}|_{11} = |A_{11} - A_{12} \cdot A_{22}^{-1} \cdot A_{21}|$ of a 2×2 -matrix $\tilde{A} = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}$; and second, consider the corresponding quasideterminant of the $k \times k$ -matrix $|\tilde{A}|_{11}$.

Basic identities. The following noncommutative analogue of Sylvester's identity is closely related with the heredity property. Let $A = (A_{ij}), i, j = 1, \dots, n$, be a matrix over a noncommutative skew-field. Suppose that the submatrix $A_0 = (a_{ij}), i, j = 1, \dots, k$, is invertible. For $p, q = k + 1, \dots, n$ set

$$b_{pq} = \left| \begin{array}{cc} A_0 & a_{1q} \\ & \vdots \\ & a_{kq} \\ a_{p1} \dots a_{pk} & a_{pq} \end{array} \right|_{pq};$$

$$B = (b_{pq}), p, q = k + 1, \dots, n.$$

Theorem 1.7. For $i, j = k + 1, \dots, n$,

$$|A|_{ij} = |B|_{ij}.$$

For example, a quasideterminant of an $n \times n$ -matrix A is equal to the corresponding quasideterminant of a 2×2 -matrix consisting of $(n - 1)$ - quasiminors of the matrix A , or to the corresponding quasideterminant of an $(n - 1) \times (n - 1)$ -matrix consisting of 2-quasiminors of the matrix A . One can use any of these procedures for a definition of quasideterminants.

II. NONCOMMUTATIVE DETERMINANTS

Let $A = (a_{ij}), i, j = 1, \dots, n$, be a generic matrix over a division algebra R , i.e. assume all square submatrices of A are invertible. For any orderings $I = (i_1, \dots, i_n)$, $J = (j_1, \dots, j_n)$ of $\{1, \dots, n\}$ define $A^{i_1 \dots i_k, j_1 \dots j_k}$ to be submatrices of A obtained by deleting rows with indices $i_1, \dots, i_k$ and columns with indices $j_1, \dots, j_k$. Then set

$$D_{I,J}(A) = |A|_{i_1 j_1} |A^{i_1 j_1}|_{i_2 j_2} |A^{i_1 i_2, j_1 j_2}|_{i_3 j_3} \dots a_{i_n j_n}.$$

In commutative case all $D_{I,J}(A)$ are equal up to a sign to the determinant of A . When A is a quantum matrix all $D_{I,J}(A)$ are equal up to a scalar to the quantum determinant of A [GR, GR1, KL]. The same is true for other well-known algebras. This gives us a reason to call the $D_{I,J}(A)$ the (I, J) -predeterminants of A . From a "categorical point of view" the expressions $D_{I,I'}$ when $I' = (i_2, i_3, \dots, i_n, i_1)$ are particularly important. Set $D_I = D_{I,I'}$. There are $n!$ expressions D_I . It is convenient to have a basic predeterminant $\Delta = D_{\{12 \dots n\}, \{12 \dots n\}}$.

We use the homological relations for quasideterminants to compare of different $D_{I,J}$'s. For example, if $I = (i_1, i_2, \dots, i_n)$, $J = (j_1, j_2, \dots, j_n)$ and $I' = (i_2, i_1, \dots, i_n)$, $J' = (j_2, j_1, \dots, j_n)$ it is enough to compare $|A|_{i_1 j_1} |A^{i_1 j_1}|_{i_2 j_2}$ and

$|A|_{i_2 j_2} |A^{i_2 j_2}|_{i_1 j_2}$. The homological relations show that these expressions are in a sense “conjugate”:

$$|A|_{i_2 j_2} |A^{i_2 j_2}|_{i_1 j_1} = |A^{i_1 j_1}|_{i_2 j_2} |A^{i_2 j_1}|_{i_1 j_2}^{-1} |A|_{i_1 j_1} |A^{i_1 j_1}|_{i_2 j_2}^{-1} |A^{i_2 j_1}|_{i_1 j_2} |A^{i_2 j_2}|_{i_1 j_1}.$$

Let R^* be the monoid of invertible elements in R and $\pi : R^* \rightarrow R^*/[R^*, R^*]$ be the canonical homomorphism of R^* .

Proposition 2.1. $\pi(\Delta(A))$ is the Dieudonne determinant of A and $\pi(D_{I,J}) = p(I)p(J)\Delta$ where $p(I)$ is the parity of ordering I .

Note, that in [Dr] P. Draxl introduced a Dieudonne predeterminant, denoted $\delta\epsilon\tau$. For a generic matrix A over a division algebra there exists the Gauss decomposition $A = UDL$ where U, D, L are upper-triangular, diagonal, and low-triangular matrices. Draxl denoted the product of diagonal elements in D from top to the bottom by $\delta\epsilon\tau(A)$ and called it the Dieudonne predeterminant of A . For non-generic matrices Draxl used the Bruhat decomposition instead of Gauss decomposition.

Proposition 2.2. $\delta\epsilon\tau(A) = \Delta(A)$.

Proof. Let $y_1, \dots, y_n$ be the diagonal elements in D from top to the bottom. As shown in [GR1], $y_k = |A^{12\dots k-1, 12\dots k-1}|_{kk}$. Then $\delta\epsilon\tau(A) = y_1 y_2 \dots y_n = \Delta(A)$.

III. QUATERNIONIC QUASIDETERMINANTS

3.1 Norms of quaternionic matrices.

Let $M(n, \mathbb{H})$ be the $\mathbb{R}$ -algebra of quaternionic matrices of order n . There exists a multiplicative functional $\nu : M(n, \mathbb{H}) \rightarrow \mathbb{R}_{\geq 0}$ such that

- i) $\nu(A) = \nu(a)$ if $A \in M(1, \mathbb{H})$, $A = (a)$,
- ii) $\nu(A) = 0$ if and only if the matrix A is non-invertible,
- iii) If A' is obtained from A by adding a left-multiple of a row to another row or a right-multiple of a column to another column, then $\nu(A') = \nu(A)$.

We will show later that for a generic quaternionic matrix $A = (a_{ij})$, $i, j = 1, \dots, n$, a norm can be defined by induction on n by formula $\nu(A) = \nu(|A|_{11})\nu(A^{11})$ and that $\nu(A)$ is a polynomial in a_{ij} 's, $\bar{a}_{ij}$'s.

3.2. Quasideterminants of quaternionic matrices.

Let $A = (a_{ij})$, $i, j = 1, \dots, n$, be a quaternionic matrix. Let I and J be two ordered sets of natural numbers $1 \leq i_1, i_2, \dots, i_k \leq n$ and $1 \leq j_1, j_2, \dots, j_k \leq n$ such that the i_p 's are distinct and the j_p 's are distinct. If $k = 1$ set $m_{I,J}(A) = a_{i_1 j_1}$. If $k \geq 2$ set

$$m_{I,J}(A) = a_{i_1 j_2} \bar{a}_{i_2 j_2} a_{i_2 j_3} \bar{a}_{i_3 j_3} a_{i_3 j_4} \dots \bar{a}_{i_k j_k} a_{i_k j_1}.$$

If the matrix A is Hermitian then

$$m_{I,J}(A) = a_{i_1 j_2} a_{j_2 i_2} a_{i_2 j_3} a_{j_3 i_3} a_{i_3 j_4} \dots a_{j_k i_k} a_{i_k j_1}.$$

Remark. The monomials $m_{I,J}(A)$ are conveniently described in terms of paths in a graph. Let Γ_n be the complete oriented 2-graph with vertices $1, \dots, n$, the arrows from p to q being labelled by a_{pq} and $\bar{a}_{qp}$. The monomial $m_{I,J}(A)$ corresponds then to a path from i_1 to j_1 .

To a quaternionic matrix $A = (a_{pq})$, $p, q = 1, \dots, n$, and to a fixed row index i and a column index j we associate a polynomial in a_{pq} 's, $\bar{a}_{pq}$'s which we call the (i, j) -th double permanent of A .

Definition 2.1. *The (i, j) -th double permanent of A is the sum*

$$\pi_{ij}(A) = \sum m_{I,J}(A),$$

taken over all orderings $I = \{i_1, \dots, i_n\}$, $J = \{j_1, \dots, j_n\}$ of $\{1, \dots, n\}$ such that $i_1 = i$ and $j_1 = j$.

Example 3.2. For $n = 2$

$$\pi_{11}(A) = a_{12}\bar{a}_{22}a_{21}.$$

For $n = 3$

$$\pi_{11}(A) = a_{12}\bar{a}_{32}a_{33}\bar{a}_{23}a_{21} + a_{12}\bar{a}_{22}a_{23}\bar{a}_{33}a_{31} + a_{13}\bar{a}_{33}a_{32}\bar{a}_{22}a_{21} + a_{13}\bar{a}_{23}a_{22}\bar{a}_{32}a_{31}.$$

For any submatrix B of A denote by B^c its complimentary submatrix. The matrix B^c is obtained from A by omitting all rows and columns containing elements from B . If B is a $k \times k$ matrix then B^c is a $(n - k) \times (n - k)$ matrix.

One of our main results is the following theorem.

Theorem 3.3. *If the quasideterminant $|A|_{ij}$ of a quaternionic matrix is defined then*

$$(3.1) \quad \nu(A^{ij})|A|_{ij} = \sum (-1)^{K(B)-1} \nu(B^c) \pi_{ij}(B)$$

where the sum is taken of all square submatrices B of A containing a_{ij} and $K(B)$ means the order of B .

Note that the quasideterminant $|A|_{ij}$ is defined if the matrix A^{ij} is invertible. In this case $\nu(A^{ij})$ is invertible and formula (3.1) gives an expression for $|A|_{ij}$.

Note that both sides in formula (3.1) are polynomials in the a_{ij} 's, the $\bar{a}_{ij}$'s with real coefficients. The right-hand side in (2.1) is a linear combination of monomials of lengths $1, 3, \dots, 2n - 1$. Let $\mu(n)$ be the number of such monomials for a matrix of order n . Then $\mu(n) = 1 + (n - 1)^2 \mu(n - 1)$.

Example 3.4. For $n = 2$

$$\nu(a_{22})|A|_{11} = \nu(a_{22})a_{11} - a_{12}\bar{a}_{22}a_{21}.$$

For $n = 3$

$$\begin{aligned} \nu(A^{11})|A|_{11} = & \nu(A^{11})a_{11} - \\ & -\nu(a_{33})a_{12}\bar{a}_{22}a_{21} - \nu(a_{23})a_{12}\bar{a}_{32}a_{31} - \nu(a_{32})a_{13}\bar{a}_{23}a_{21} - \nu(a_{22})a_{13}\bar{a}_{33}a_{31} + \\ & + a_{12}\bar{a}_{32}a_{33}\bar{a}_{23}a_{21} + a_{12}\bar{a}_{22}a_{23}\bar{a}_{33}a_{31} + a_{13}\bar{a}_{33}a_{32}\bar{a}_{22}a_{21} + a_{13}\bar{a}_{23}a_{22}\bar{a}_{32}a_{31}. \end{aligned}$$

Example 3.4 shows how to simplify the general formula for quasideterminants of matrix of order 3 (see example 1.2) for quaternionic matrices.

Formula (3.1) gives a recurrence relation for norms of quaternionic matrices. From this relation one can obtain an expression of $\nu(A)$ as a polynomial in a_{ij} 's and their conjugates (see also subsection 4.3 below).

Corollary 3.5.

$$\nu(A)\nu(A^{ij}) = \sum (-1)^{K(B)-1} \nu(B^c) \pi_{ij}(B) \left(\sum (-1)^{K(B)-1} \nu(B^c) \overline{\pi_{ij}(B)} \right).$$

Example 3.6. For $n = 2$

$$\nu(A) = \nu(a_{11})\nu(a_{22}) + \nu(a_{12})\nu(a_{21}) - a_{12}\bar{a}_{22}a_{21}\bar{a}_{11} - a_{11}\bar{a}_{21}a_{22}\bar{a}_{12}.$$

For a square quaternionic matrix $A = (a_{ij})$ set $Q_{ij}(A) = |A|_{ij}\nu(A^{ij})$. Then one can rewrite formula (3.1) as

$$Q_{ij}(A) = \nu(A^{ij})a_{ij} - \sum_{p \neq i, q \neq j} a_{iq} \overline{Q_{pq}(A^{ij})} a_{pj}.$$

If A is a Hermitian matrix then $\bar{Q}_{pq}(A^{ij}) = Q_{qp}(A^{ij})$ and

$$Q_{ij}(A) = \nu(A^{ij})a_{ij} - \sum_{p \neq i, q \neq j} a_{iq} Q_{qp}(A^{ij}) a_{pj}.$$

When A is a complex matrix, formula (3.1) can be written as

$$\det A \det \bar{A}^{ij} = (-1)^{i+j} (a_{ij} \det \bar{A}^{ij} - \sum_{p \neq i, q \neq j} a_{iq} \det \bar{A}^{ij} \det A^{ip,jq} a_{pj}).$$

This implies the standard row/column expansion of the determinant.

IV. DETERMINANTS OF QUATERNIONIC MATRICES

4.1. Dieudonne determinant.

Let $A = (a_{ij})$, $i, j = 1, \dots, n$ be a generic quaternionic matrix. Recall that for any orderings $I = (i_1, \dots, i_n)$, $J = (j_1, \dots, j_n)$ of $\{1, \dots, n\}$ we defined expressions

$$D_{I,J}(A) = |A|_{i_1 j_1} |A^{i_1 j_1}|_{i_2 j_2} |A^{i_1 i_2, j_1 j_2}|_{i_3 j_3} \dots a_{i_n j_n}.$$

In the quaternionic case the Dieudonne determinant D maps

$$D : M_n(\mathbb{H}) \rightarrow \mathbb{R}_{\geq 0}$$

(see [As]).

The following proposition generalizes a result from [VP].

Proposition 4.1. *In the quaternionic case $|D_{I,J}(A)|$, the absolute value of $D_{I,J}(A)$, is equal to $D(A)$ for any I, J .*

A proof of Proposition 4.1 follows from the homological relations for quasideterminants.

4.2. Moore determinants of Hermitian quaternionic matrices.

A quaternionic matrix $A = (a_{ij})$, $i, j = 1, \dots, n$ is Hermitian if $a_{ji} = \bar{a}_{ji}$ for all i, j . It follows that all diagonal elements of A are real numbers and that the submatrices A^{11} , $A^{12,12}$, $\dots$ are Hermitian.

A determinant of Hermitian matrices was introduced by E.M. Moore in 1922 [M, MB]. Here we recall the original definition.

Let $A = (a_{ij})$, $i, j = 1, \dots, n$, be a matrix over a ring. Let σ be a permutation of $\{1, \dots, n\}$. Write σ as a product of disjoint cycles such that each cycle starts with a smallest number. Since disjoint cycles commute, we may write

$$\sigma = (k_{11} \dots k_{1j_1})(k_{21} \dots k_{2j_2}) \dots (k_{m1} \dots k_{mj_m})$$

where for each i , we have $k_{i1} < k_{ij}$ for all $j > 1$, and $k_{11} > k_{21} \dots > k_{m1}$. This expression is unique. Let $p(\sigma)$ be a parity of σ . Define Moore determinant $M(A)$ by

$$(4.1) \quad M(A) = \sum_{\sigma \in S_n} p(\sigma) a_{k_{11}, k_{12}} \dots a_{k_{1j_1}, k_{11}} a_{k_{21}, k_{22}} \dots a_{k_{mj_m}, k_{m1}}.$$

One can write cycles starting with the largest element. This will lead to the same result.

If A is Hermitian quaternionic matrix then $M(A)$ is a real number. Moore determinants have nice features and are widely used (see, for example, [Al, As, Dy1]).

The following Theorem 4.3 shows that determinants of Hermitian quaternionic matrices can be obtained through our general approach. We prove first, that for quaternionic Hermitian matrix A determinants $D_{I,I}(A)$ coincide up to a sign.

Recall that $\Delta(A) = D_{I,I}(A)$ for $I = \{1, \dots, n\}$ and that $\Delta(A)$ is a pre-Dieudonné determinant in the sense of [Dr]. If A is Hermitian then $\Delta(A)$ is a product of real numbers and, therefore, $\Delta(A)$ is real.

Proposition 4.2. *Let $p(I)$ be the parity of the ordering I . Then $\Delta(A) = p(I)p(J)D_{I,J}(A)$.*

The proof follows from homological relations for quasideterminants.

Theorem 4.3. *Let A be a Hermitian quaternionic matrix. The $\Delta(A) = M(A)$.*

Proof. We use Sylvester formula for quasideterminants (Theorem 1.7).

For $i, j = 2, \dots, n$ set

$$B_{ij} = \begin{pmatrix} a_{11} & a_{1j} \\ a_{i1} & a_{ij} \end{pmatrix}.$$

B_{ij} is a Hermitian matrix. Let $b_{ij} = M(B_{ij})$ and $c_{ij} = |B_{ij}|_{11}$.

Note that $B = (b_{ij})$ and $C = (c_{ij})$ also are Hermitian matrices. It follows from (3.1) that $M(A) = a_{nn}^{2-n} M(B)$. Note, that $M(B) = a_{nn}^{n-1} M(C)$, therefore, $M(A) = a_{nn} M(C)$.

By the noncommutative Sylvester formula (Theorem 1.7), $|A|_{11} = |C|_{11}$, $|A^{11}|_{22} = |C^{11}|_{22}$, So,

$$|A^{11}|_{22} |A^{11}|_{22} \dots |A^{12\dots n-1, 12\dots n-1}|_{n-1, n-1} = |C^{11}|_{22} |C^{11}|_{22} \dots |C^{12\dots n-1, 12\dots n-1}|_{n-1, n-1}.$$

The left-hand product equals $\Delta(A)a_{nn}^{-1}$ and the right-hand product equals $\Delta(C)$, so $\Delta(A) = \Delta(C)a_{nn} = M(A)$.

4.3. Moore determinants and norms of quaternionic matrices.

For a generic quaternionic matrix $A = (a_{ij})$, $i, j = 1, \dots, n$, define quaternionic norm $\nu(A)$ by induction:

- i) $\nu(A) = \nu(a)$ if A is the 1×1 -matrix (a) ,
- ii) $\nu(A) = \nu(|A|_{11})\nu(A^{11})$.

It follows that $\nu(A) \in \mathbb{R}_{\geq 0}$.

Proposition 4.4. *For generic matrices A, B*

- i) $\nu(AB) = \nu(A)\nu(B)$,
- ii) $\nu(A) = \Delta(A)\Delta(A^*) = \Delta(AA^*)$.

Our proof is based on properties of quasideterminants.

Since AA^* is a Hermitian matrix one has the following

Corollary 4.5. $\nu(A) = M(AA^*)$.

It follows that $\nu(A)$ is a polynomial in the a_{ij} 's and the $\bar{a}_{ij}$'s and so it can be defined for all matrices A by continuity. The continuation possesses the multiplicative property.

4.4. Study determinants.

In Section 1.1 a homomorphism $\theta : \mathbb{H} \rightarrow M(2, \mathbb{C})$ was defined. This homomorphism can be extended to homomorphism of matrix algebras

$$\theta_n : M(n, \mathbb{H}) \rightarrow M(2n, \mathbb{C}).$$

Let $A = (a_{ij}) \in M(n, \mathbb{H})$, set $\theta_n(A) = (\theta(a_{ij}))$.

In 1920, E. Study [S] defined a determinant $S(A)$ of a quaternionic matrix A of order n by setting $S(A) = \det \theta_n(A)$. Here $\det$ is the standard determinant of a complex matrix.

The following proposition is well known (see [As]).

Proposition 4.6. *For any quaternionic matrix A*

$$S(A) = M(AA^*).$$

The proof in [As] was based on properties of eigenvalues of quasideterminant matrices. Our proof based on Sylvester identity and homological relations actually shows that $S(A) = \nu(A)$ for a generic matrix A .

V. ALGEBRAS WITH ANTI-INVOLUTIONS

All our results are valid for algebra R with anti-involutions $I : R \rightarrow R$ such that subalgebra $R_0 = \{r \in R : I(r) = r\}$ is central and the monoid of invertible elements in R is open in some topology.

Let R be an algebra over a field F . An anti-involution of R is a F -linear map $I : R \rightarrow R$ such that $I(xy) = I(y)I(x)$ and $I^2(x) = x$ for any $x, y \in R$. It is common to write $\bar{x}$ instead of $I(x)$ for $x \in R$.

REFERENCES

- [Ad] S.L. Adler, *Quaternionic quantum mechanics and quantum fields*, 1995.
- [Al] S. Alesker, *Non-commutative linear algebra and plurisubharmonic functions of quaternionic variables*, arXiv:math.CV/010429 (2002).
- [As] H. Aslaksen, *Quaternionic determinants*, Math. Intelligencer . **15** (1996), no. 3, 57-65.
- [B] F. A. Berezin, *Introduction to Super analysis*, 1987.
- [Ca] A. Capelli, *Über die Zurückführung der Cayley'schen Operationen Ω und gewöhnliche Polaroperationen*, Math. Ann. . **29** (1887), 331-338.
- [C] A. Cayley, *On certain results relating to quaternions*, Collected math papers **1** (1989), 123-126.

- [Di] J. Dieudonne, *Les determinantes sur un corps non commutatif*, Bull. Soc. Math. France . **71** (1943), 27-45.
- [Dr] P. Draxl, *Skew fields*, London Math. Soc. Lecture notes **81** (1983).
- [Dy1] F. Dyson, *Quaternion determinants*, Helv. Phys. Acta **45** (1972), 289-302.
- [EGR] P. Etingof, I. Gelfand, and V. Retakh, *Factorization of Differential Operators, Quasideterminants, and Nonabelian Toda Field Equations*, Math. Res. Letters **4** (1997), no. 2-3, 413-425.
- [EGR1] P. Etingof, I. Gelfand, and V. Retakh, *Nonabelian Integrable Systems, Quasideterminants, and Marchenko Lemma*, Math. Res. Letters **5** (1998), no. 1, 1-12.
- [ER] P. Etingof and V. Retakh, *Quantum Determinants and Quasideterminants*, Asian J. of Mathematics **3** (1999), no. 2, 345-352.
- [F] D. Foata, *A noncommutative version of the matrix inversion formula*, Adv. in Math. . **31** (1996), no. 3, 330-349.
- [GKLLRT] I. Gelfand, D. Krob, A. Lascoux, B. Leclerc, V. Retakh, and J.-Y. Thibon, *Noncommutative Symmetric Functions*, Advances in Math **112** (1995), no. 2, 218-348.
- [GR] I. Gelfand and V. Retakh, *Determinants of Matrices over Noncommutative Rings*, Funct. Anal. Appl. **25** (1991), no. 2, 91-102.
- [GR1] I. Gelfand and V. Retakh, *A Theory of Noncommutative Determinants and Characteristic Functions of Graphs*, Funct. Anal. Appl. **26** (1992), no. 4, 1-20.
- [GR2] I. Gelfand and V. Retakh, *A Theory of Noncommutative Determinants and Characteristic Functions of Graphs. I*, Publ. LACIM, UQAM, Montreal **14** (1993), 1-26.
- [GR3] I. Gelfand and V. Retakh, *Noncommutative Vieta Theorem and Symmetric Functions*, Gelfand Mathematical Seminars 1993-95, Birkhauser, Boston (1996), 93-100.
- [GR4] I. Gelfand and V. Retakh, *Quasideterminants, I*, Selecta Math **3** (1997), no. 4, 517-546.
- [GV] V. Goncharenko and A. Veselov, *Darboux Transformations for Multidimensional Schrödinger Operators*, J. Phys. A: Math. Gen. **31** (1998), 5315-5326.
- [H] A. Heyting, *Die Theorie der linearen Gleichungen in einer Zahlenspezies mit nichtkommutativer Multiplikation*, Math. Ann . **98** (1927), 465-490.
- [J] C. Joly, *Supplement to Oeuvres of Hamilton*, 1900.
- [KL] D. Krob and B. Leclerc, *Minor Identities for Quasi-Determinants and Quantum Determinants*, Comm. Math. Phys. **169** (1995), no. 1, 1-23.
- [KS] P. P. Kulish, E. K. Sklyanin, *Quantum spectral transform method. Recent developments*, Math. Lecture Notes in Physics . **151** (1982), no. 3, 61-119.
- [LST] B. Leclerc, T. Scharf, and J.-Y. Thibon, *Noncommutative Cyclic Characters of Symmetric Groups*, J. of Combinatorial Theory, Series A **75** (1996), 55-69.
- [M] E.H. Moore, *On the determinant of an hermitian matrix of quaternionic elements*, Bull. Amer. Math. Soc. **28** (1922), 161-162.
- [MB] E.H. Moore and R.W. Barnard, *General Analysis*, 1935.
- [Mo] A. Molev, *Noncommutative Symmetric Functions and Laplace Operators for Classical Lie Algebras*, Lett. Math. Phys. **35** (1995), no. 2, 135-143.
- [Mo1] A. Molev, *Gelfand-Tsetlin Bases for Representations of Yangians*, Lett. Math. Phys. **30** (1994), no. 2, 53-60.
- [O] O. Ore, *Linear equations in noncommutative rings*, Ann. Math. . **32** (1996), 463-477.
- [P] A. Polischuk, *Triple Massey Products on Curves, Fay's Trisecant Identity and Tangents to the Canonical Embedding*, preprint math.AG/0109007 (2001).
- [R] A. R. Richardson, *Hypercomplex determinants*, Messenger of Math. . **55** (1926), 145-152.
- [RS] A. Razumov, M. Saveliev, *Maximally Nonabelian Toda Systems*, Nuclear Physics B **494** (1997), no. 3.

- [RRV] V. Retakh, C. Reutenauer, and A. Vaintrob, *Noncommutative Rational Functions and Farber's Invariants of Boundary Links*, in: Differential Topology, Infinite-dimensional Lie Algebras and Applications. Amer Math Soc Transl. **194** (1999), 237-246.
- [RSh] V. Retakh and V. Shander, *Schwarz Derivative for Noncommutative Differential Algebras*, Advances in Soviet Math. **17** (1993), 139-154.
- [Re] C. Reutenauer, *Inversion Height in Free Fields*, Selecta Mathematica **2** (1996), no. 1.
- [SK] M. Sato, M. Kashiwara, *The determinant of matrices of pseudo-differential operators*, Proc. Japan Acad. of Sci. . **51** (1975), no. 1, 17-19.
- [S] E. Study, *Zur Theorie der Lineare Gleichungen*, Acta Math. **42** (1920), 1-61.
- [Sch] M. Schork, *The bc-sustem of higher rank revisited*, J. Phys. A: Math. Gen. **35** (2001), 2627-2637.
- [VP] P. Van Praag, *Sur le norme reduite du determinant de Dieudonne des matrices quaterniennes*, J. Algebra **136** (1991), 265-274.
- [W] R. F. Williams, *A new zeta function, natural for links*, From Topology to Computation (Smalefest volume) (1993), 270-278.
- [Z] F. Zhang, *Quaternions and matrices of quaternions*, Linear Algebra Appl. **251** (1997), 21-57.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854

E-mail address:

I. G.: `igelfand@math.rutgers.edu`

V. R.: `vretakh@math.rutgers.edu`

R. W.: `rwilson@math.rutgers.edu`